



RESEARCH ARTICLE

Open Access

The role of FGF-2 and BMP-2 in regulation of gene induction, cell proliferation and mineralization

Millie Hughes-Fulford^{1,2,3,4*}, Chai-Fei Li⁴

Abstract

Introduction: The difficulty in re-growing and mineralizing new bone after severe fracture can result in loss of ambulation or limb. Here we describe the sequential roles of FGF-2 in inducing gene expression, cell growth and BMP-2 in gene expression and mineralization of bone.

Materials and methods: The regulation of gene expression was determined using real-time RTPCR (qRTPCR) and cell proliferation was measured by thymidine incorporation or fluorescent analysis of DNA content in MC3T3E1 osteoblast-like cells. Photomicroscopy was used to identify newly mineralized tissue and fluorescence was used to quantify mineralization.

Results: Fibroblast growth factor-2 (FGF-2) had the greatest ability to induce proliferation after 24 hours of treatment when compared to transforming growth factor beta (TGFβ), insulin-like growth factor-1 (IGF-1), bone morphogenic protein (BMP-2), platelet derived growth factor (PDGF) or prostaglandin E₂ (PGE₂). We found that FGF-2 caused the most significant induction of expression of early growth response-1 (*egr-1*), *fgf-2*, *cyclo-oxygenase-2 (cox-2)*, *tgfβ* and matrix metalloproteinase-3 (*mmp-3*) associated with proliferation and expression of angiogenic genes like vascular endothelial growth factor A (*vegfa*) and its receptor *vegfr1*. We found that FGF-2 significantly reduced gene expression associated with mineralization, e.g. *collagen type-1 (col1a1)*, *fibronectin (fn)*, *osteocalcin (oc)*, *IGF-1*, *noggin*, *bone morphogenic protein (bmp-2)* and *alkaline phosphatase (alp)*. In contrast, BMP-2 significantly stimulated expression of the mineralization associated genes but had little or no effect on gene expression associated with growth.

Conclusions: The ability of FGF-2 to re-program a mineralizing gene expression profile to one of proliferation suggests that FGF-2 plays a critical role of osteoblast growth in early fracture repair while BMP-2 is instrumental in stimulating mineralization.

Introduction

The mechanisms that regulate bone growth and mineralization remain poorly understood. The cellular events involved in bone formation include chemotaxis of osteoblast precursors, growth factor (GF) production, proliferation of committed osteoblast precursors, and the differentiation (mineralization) of osteoblasts. Bone formation requires expression of structural proteins such as collagen type I, osteocalcin, noggin and runx2 during mineralization [1]. Numerous studies suggest that a

variety of growth factors such as FGF-2, TGFβ, IGF-1, PDGF and PGE₂ act as autocrine and paracrine hormones to regulate bone cell proliferation [2]. FGF-2 is an important modulator of bone formation *in vitro* and *in vivo* [3,4]. FGF-2 is tightly bound to the bone matrix and can be extracted as a biologically active GF [5] and is thought to play a major role in wound healing [6,7].

To evaluate the physiological activity of FGF-2 and other growth factors, we studied their relative ability to influence proliferation of osteoblasts at a site of injury in a mineralized culture. MC3T3-E1 is a cloned mouse osteoblast-like cell line that retains synthetic functions of bone. When treated with differentiation media, these cultured osteoblasts have the ability to differentiate,

* Correspondence: millie.hughes-fulford@ucsf.edu

¹Department of Research, Veterans Affairs Medical Center, 4150 Clement Street, San Francisco, CA 94121, USA

Full list of author information is available at the end of the article

including synthesis of alkaline phosphatase [8], type I collagen [9], osteocalcin [10,11] and mineralized matrix containing hydroxyapatite crystals [12].

We have previously reported that FGF-2 is induced by mechanical stress [13,14] and causes proliferation after mechanical stress. FGF-2 is an immediate-early gene that is regulated by both PKA and MAPK signal transduction pathways [15]. Here we report that FGF-2 induces expression of growth-related genes and down-regulates genes responsible for differentiation and mineralization. In addition, BMP-2 is considerably more effective than FGF-2 in inducing new mineralization.

Materials and methods

Materials

We obtained GFs from Amgen, Thousand Oaks, CA. FGF-2 and IGF-1 from R & D Systems, Minneapolis, MN. TGF β , PDGF and dmPGE₂ are from Cayman Chemical, Ann Arbor, Michigan. Cell culture supplies (α MEM, fetal calf serum, trypsin and antibiotics) were obtained through the tissue culture facility at the University of California, San Francisco. Cell culture dishes were purchased from Corning, Corning, New York. Rhodamine-phalloidin is from Invitrogen, Carlsbad, California. Tritiated thymidine and ³⁵S methionine are from Amersham, Arlington Heights, IL. All other materials came from standard laboratory suppliers. MC3T3E1 osteoblast-like cells, a cloned cell line, established by Kodama [8,12] were used in this study at early passage number.

Methods

We maintained cloned MC3T3-E1 osteoblast-like cells in normal media (NM) consisting of alpha MEM medium with 10% fetal calf serum (FCS), 1% antibiotic solution and 1% glutamine solution and subcultured the cells every 3 to 4 days. The cells were subcultured by incubating with trypsin for five minutes and resuspending at a concentration of 3×10^5 cells/ml. For experiments, we grew the cells in the NM above, using multi-well plates. After three days, the cells reach confluence and mineralization medium (MM) was added. MM is alpha MEM medium with 5% fetal calf serum (FCS), 1% antibiotic solution and 1% glutamine solution supplemented with ascorbic acid (50 μ g/ml) and β -glycerol phosphate (10 mM) to support mineralization. The cultures were then incubated for 1-2 more days for mineralization studies. We used at least triplicate independent biological samples in multiple experiments for data collection.

Protein Assay

Protein concentration was determined by Bio-Rad DC protein assay (Bio-Rad, CA) according to manufacturer's protocol.

Microscopy

At the conclusion of the 24 or 48 hour incubation, the coverslip was removed. The specimen was rinsed five times in room temperature phosphate buffered saline (PBS) and fixed. We then visualized the mineralizing cells with 2% Alizarin Red. After rinsing in distilled water and air drying the samples, we mounted the coverslips on microscope slides using Fluoromount and examined and photographed the cells on a Zeiss Axioskop using 20 \times .

Tritiated thymidine incorporation into DNA

At the conclusion of the 24 hour incubation, the culture medium was removed and the cells were incubated for 15 minutes at 37 $^{\circ}$ C in 1 ml PBS containing tritiated thymidine (4 μ Ci/ml) as described previously [16]. Following this incubation, the PBS was removed and the cells were washed 3 times with ice cold trichloroacetic acid (TCA) followed by ice cold ethanol and allowed to air dry. Then 1 ml of sarkosyl lysing buffer was added to each well; all the cells were solubilized after 30 minutes. Finally, after mixing the resulting solution with a pipette, radioactivity was counted in a scintillation counter and protein content was measured. The data was calculated and expressed as disintegrations per minute (DPM) per microgram protein.

Alizarin Red visualization of mineralization

Alizarin Red (2%) stained cells were incubated with 10% acetic acid for 30 minutes to release bound Alizarin Red into solution. The solution was neutralized with 10% ammonium hydroxide and the absorbance of Alizarin Red was measured at 450 nm using a microplate reader. Data is expressed in absolute amounts according to a standard curve.

RNA Isolation

RNA were isolated through the use of the RNeasyTM Mini kit (QIAGEN, Valencia, CA) or TriReagentTM according to the manufacturer's protocol. For RNeasyTM Mini kit RNA isolation, cells were seeded in 6-well plates with α MEM media supplemented with 10% FCS, then downregulated and activated as indicated in the figure legends. Cells were lysed using 350 μ l of buffer RLT (supplied in kit) containing 2-mercaptoethanol (Biorad, Hercules, CA). The lysate was then placed into QIAshredder homogenizer (QIAGEN, Valencia, CA) and centrifuged at 20,000 rpm for 2 minutes. 350 μ l of 70% ethanol was added to the flow through, mixed, and centrifuged in the RNeasyTM Mini column (supplied in kit) for 15 s at 20,000 rpm. Flow through was discarded and the column was washed with 700 μ l of buffer RW1 (supplied in kit) for 15 s at 20,000 rpm. Two additional washes were performed with 500 μ l of buffer RPE

(supplied in kit) at 20,000 rpm for 15 s and 2 minutes, respectively. The flow through was discarded and the column placed in a sterile 1.5 ml collection tube. Depending on the expected yield, 20-50 µl RNase-free water is pipetted into the column and centrifuged for 1 minute at 20,000 rpm. The samples are then stored at -80°C until further analysis.

Reverse Transcription (RT)

1.5 µg of RNA was added to 30 µl reverse transcriptase (RT) reaction buffer containing 5 mM MgCl₂, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1 mM dNTPs, 2.5 µM oligo d(T) primer, 2.5 U/µl of MuLV, and 1 U/µl of RNase inhibitor. The RT reaction was incubated at room temperature for 10 min, 42°C for 30 min, inactivated at 99°C for 5 min, and cooled at 5°C for 5 min.

Real-time Quantitative RT-PCR Reaction (qRTPCR)

2 µl of cDNA from the RT reaction was added to 20 µl real-time quantitative polymerase chain reaction (qPCR) mixture containing 10 µl of 2× SYBR® Green PCR Master Mix (Applied Biosystems, Foster City, CA) and 12 pmol oligonucleotide primers. PCRs were carried out in a Bio-Rad MyiQ Single-Color Real-Time PCR Detection System (Bio-Rad, Hercules, CA). The thermal profile was 50°C for 2 min, 95°C for 10 min to activate the Taq polymerase, followed by 50 amplification cycles, consisting of denaturation at 95°C for 1 min 40 s, annealing at 63°C for 1 min 10 s and elongation at 72°C for 1 min 40 s. Fluorescence was measured and used for quantitative purposes. At the end of the amplification period, melting curve analysis was performed to confirm the specificity of the amplicon. RNA samples were normalized to *cyclophilin* (*CPhi*) internal standard. Relative quantification of gene expression was calculated by using $2^{-(C_t \text{ gene T} - C_t \text{ CPhi T}) - (C_t \text{ gene 0 hr} - C_t \text{ CPhi 0 hr})}$ equation, where “C_t gene T” represents the calculated threshold cycle (C_t) of a time point of each sample other than 0 hr, or each treatment other than control. Relative gene absolute abundance was calculated using $2^{\text{sup}> (C_t \text{ gene T} - C_t \text{ CPhi T})}$ as previously described [17] allows us to compare the abundance of the gene between other genes and experiments. The resulting numbers were then multiplied by 10,000 for better graphical presentation. Primer sequence information was previously published [18-22]. All data derived using qRTPCR was from multiple experiments with at least triplicate independent biological samples.

Results

Growth factor effect on cell proliferation DNA synthesis

As seen in Table 1, in the absence of any added compounds there were small and unremarkable changes in DNA synthesis with IGF-1 and PDGF; in contrast,

Table 1 Effect of growth factors on protein synthesis in wounded mineralized osteoblasts

Treatment	Thymidine incorporation DPM × 10 ³ /ug protein
con	37.6 ± 2.9
IGF-1	42.3 ± 4.2
FGF-2	114.3 ± 11
TGFβ	65.2 ± 12
PDGF	39.8 ± 7.2
BMP-2	41.5 ± 5.6
PGE2	84.1 ± 23.1

Representative experiment showing the effects of IGF-1 (20 ng/ml), FGF-2 (2.0 ng/ml), TGFβ (2 ng/ml), PDGF (3 ng/ml), BMP-2 (100 ng), PGE₂ (2 µg/ml) on proliferation/mg protein of MC3T3-E1 osteoblasts after 24 hours of treatment (n = 4).

FGF-2, TGFβ and PGE2 significantly enhanced thymidine incorporation within 24 hours of treatment. TGFβ stimulated thymidine incorporation more than 2 fold while FGF-2 and PGE2 increased DNA synthesis more than 4.5 and 3.3 fold respectively.


Regulation of FGF-2 induced gene expression

Using qRTPCR, we found that FGF-2 dramatically induced *egf-1*, *fgf-2*, *cox-2*, *tgfβ*, *mmp3*, *vegfa* and *vegfr1* over a 24 hour period each displaying a different sequential temporal pattern of gene induction (Figure 1). *Vegfa* and *vegfr1* are associated with angiogenesis while *mmp3*, is associated with increased migration. *Tgfβ*, *fgf-2*, *egr-1* and *cox-2* are key genes in regulation of osteoblast proliferation.

Interestingly, we found that FGF-2 also significantly decreased expression of other genes associated with mineralization including *col1a1*, *fn*, *bmp-2*, *oc*, *run-x*, and *noggin*. IGF-1, a known differentiation factor, was significantly decreased by FGF-2 treatment. (Figure 2).

Relative abundance of genes regulated by FGF-2 and BMP-2

Since FGF-2 increased growth associated genes, we used BMP-2, a known promoter of mineralization, to study relative abundance of gene expression in mineralizing cells after 24 hours of treatment. As seen in Table 2, we found that BMP-2 treatment caused significant increases in genes associated with mineralization including *cola1*, *fn*, *noggin* and *oc*. Moreover, BMP-2 treatment caused little or no changes in expression of genes associated with angiogenesis and migration e.g. VEGF and MMP3. When compared with relative gene abundance of FGF-2 treated cells (Figure 3) we found that in general, BMP-2 maintained the mineralizing RNA profile of *igf-1*, *alp*, and *bmp-2* and significantly increased expression of other genes associated with mineralization like *col1a1*, *fn*, *ilgf-1*, *noggin* and *oc*. Fgf-2, on the other hand, significantly suppressed expression of mineralizing genes.




Relative mineralization of FGF-2 and BMP-2 treated cells

As seen in Figure 4 and Table 3, BMP-2 treatment enhances mineralization of the cells as shown by uptake and presence of Alizarin Red after cultures were grown to confluence and then treated with BMP-2 or FGF-2 for 24 to 48 hours. Cells were then washed and stained with 2% Alizarin Red and results determined using photography or fluorescence analysis at 48 hours of treatment.

Discussion

Bone formation during injury repair is a multi-step series of events modulated by an integrated cascade of gene expression that initially supports the proliferation stage. The later mineralization stage is associated with the sequential expression of genes that support biosynthesis, organization and mineralization of the bone extracellular matrix. Mineralization requires expression of structural proteins such as collagen type I, osteocalcin, as well as noggin and runx2 which aid in mineralization [1].



Transcriptional control defines the regulatory events necessary for both stages of bone formation [23]. There is a general consensus that during injury GFs are released from the wounded bone matrix and promote healing [24]. In this study, we have documented the relative efficiency of bone growth factors FGF-2, TGF β , and PGE $_2$ markedly enhanced the synthesis of the total protein content of the dishes (Table 1)

Rate of proliferation was dependent on the specific GF. FGF-2, TGF β and PGE $_2$ significantly promote growth, with FGF-2 having the highest efficacy and the lowest dose. FGF-2 produced a distinct pattern of gene expression. FGF-2 down regulates genes associated with mineralization while it induces genes associated with proliferation and angiogenesis, a finding supported by observations of others [25]. Since *cox-2* had a 27-fold induction by FGF-2, we examined the effect of the COX-2 product, PGE $_2$ on proliferation. We found that PGE $_2$ increased DNA synthesis by 3.3 fold significantly higher than TGF β , IGF-1, PDGF, suggesting that its induction by FGF-2 helps complete the FGF-2 full induction of osteoblast growth. These data also suggest that FGF-2 may be an important regulator of migration, angiogenesis and proliferation during the first stage of healing a critical defect since it induces *mmp3*, *vegfa* and *vegfr1* expression. In data not shown, FGF-2 had no effect on expression of *mmp-1*. Moreover, FGF-2

Table 2 Relative abundance of gene expression in FGF-2 and BMP-2 treated cells

Gene	Non-treated		FGF-2 treated		BMP-2 treated		FGF-2 vs BMP-2
	Average	SD	Average	SD	Average	SD	p-value
Collagen Type I	85,081.73	2,5316.39	**678.21	358.27	*170,243.43	24,493.77	0.0003
Fibronectin	55,827.93	1,2119.18	*28,432.19	1195.92	**239,750.67	23,464.19	0.0001
IGF1	3,249.41	689.70	**50.65	13.30	4,193.34	739.19	0.0006
RUNX2	349.09	40.63	**674.95	63.04	1,043.65	783.29	n.s.
VEGFA	109.49	38.86	**5,132.66	755.22	537.13	379.66	0.0007
TGFβ	93.08	10.55	**245.40	41.93	*185.20	38.34	n.s.
ALP	58.30	34.81	13.39	11.68	91.77	23.15	0.0064
OC	16.20	3.19	**1.38	0.65	*34.04	6.11	0.0008
Noggin	7.11	2.77	*1.61	0.49	2.41	1.76	n.s.
BMP-2	0.40	0.12	**0.06	0.01	0.38	0.05	0.0004
MMP3	0.03	0.03	**4.04	0.97	0.12	0.14	0.0023

This table shows the relative abundance of gene expression in mineralizing MC3T3-E1 cells after 24 hours of treatment with FGF-2 (5 ng) or BMP-2 (100 ng). Total RNA was harvested 24 hours after the addition using Qiagen RNeasy kit. A two-step RT-qPCR was performed. Each data point represents the mean ± SD of three biological independent samples. *p < 0.05; **p < 0.01; ***p < 0.0001 against 0 hour control samples with 2 tail student t test.

induced its own message as well as TGFβ, but significantly reduced expression of BMP-2, osteocalcin, noggin, runx2, collagen type I and IGF-1, genes which are associated with mineralization.

As described by others, bone formation is divided into two phases, proliferation and mineralization [2,26-29]. These two stages are the result of a specific sequential regulation of gene expression from the early phase of osteoblast proliferation to the final steps of mineralization.

Once the cells start mineralizing, cell division and DNA synthesis dramatically slow down and eventually cease. When an injury occurs in mineralized tissue, GFs like FGF-2 are released and start a new proliferation stage to heal the defect. The increase in cell replication in a mineralizing cell likely represents a de-differentiation from the mineralizing phase to the growing phase, and increases expression of GFs most likely induce proliferation. Treatment of the mineralized defect model with FGF-2 resulted in gene expression that corresponds to de-differentiation (e.g. significant increases in growth related genes *egf-1*, *fgf-2*, *cox-2*, *TGFβ*, *vegfa*, *vegfr* and *mmp3* and down-regulation of mineralizing related genes). *Vegf* and *vegfr1* are primary regulators of angiogenesis, while *MMP3* is thought to play a major role on cell behaviors such as proliferation and migration [30] which may explain the ability of the FGF-2 to enable the cultured cells to fill the defect void efficiently. The fact that FGF-2 induces its own expression suggests that after injury, the FGF-2 released from the wound matrix could promote its own expression, making it a feed-forward loop.

Although Figures 1 and 2 demonstrate the relative FGF-2 regulation and sequential expression of growth, angiogenic and chemotactic genes and depresses expression of mineralization-related genes, these figures do not tell us the relative abundance of the genes. In Table 2, we determined the relative abundance of genes in three groups after 24 hours; with or without treatment with FGF-2 or BMP-2. FGF-2 caused a significant increase in abundance of genes associated with proliferation, chemotaxis and angiogenesis. Moreover, the addition of FGF-2 to the mineralized wounded cultures caused a marked decrease in abundance of *coll1a1* as well as *fn*, *igf-1*, *noggin*, *oc*, *bmp-2* and *alp* message. In the early stages of mineralization, the major protein (greater than 20%) synthesized by the osteoblast is collagen, however





Figure 3 FGF-2 and BMP-2, the yin and yang of mineralization: Contrast of effect of 24 hours of treatment with FGF-2 or BMP-2 on fold increase in abundance of mineralization-related gene expression. Mineralizing MC3T3-E1 cells were prepared as described in Materials and Methods. They were then treated with either FGF-2 or BMP-2 for 24 hours at which time RNA was collected and analyzed for relative abundance using qRT-PCR. Each bar represents mean ± SD triplicate independent biological samples each time point corrected to cyclophilin. (*p < 0.05; **p < 0.01 with two-tail student t-test compare to 0 hour of each gene). * < 0.05; ** < 0.01; *** < 0.0001.



collagen is not a major component of the proliferating cell, suggesting that FGF-2 stimulates proliferation partly through its ability to drastically reduce the relative abundance of a majority of the mineralizing-associated genes. The dramatic reduction of IGF-1 by FGF-2 suggests a role for IGF-1 in mineralization, this is in agreement with findings of others that demonstrated IGF-1 to be a major factor in bone mineralization [31-33] using the IGF-1 null mouse. In contrast, in cells treated with BMP-2, the relative abundance of *col1a1*, *fn*, *oc*, and *tgfb3* were dramatically induced while BMP-2 had no

significant effect on genes related to growth, angiogenesis or chemotaxis. These data suggest that BMP-2 may be the best GF to use for the mineralization stage but not the proliferation stage of bone formation. This finding may help explain studies by others [34] who discovered that a *delayed* administration of BMP-2 to a fracture resulted in better repair of critical size defects. It is likely that the delay of BMP-2 treatment allowed a longer period of proliferation prior to BMP-2 promotion of mineralization. Our findings in Table 2, 3 and Figure 3 support the hypothesis that FGF-2 and BMP-2 are required at different stages of bone repair.

Table 3 Mineralization of cells with BMP-2

Treatment	Relative abundance
NM	5.6 ± 1.7
NM + 5 ng/ml FGF-2	5.3 ± .1
NM + 50 ng/ml BMP-2	16.2 ± 4.2
MM	9.1 ± 2.0
MM + 5 ng/ml FGF-2	4.9 ± 1.1
MM + 50 ng/ml BMP-2	55.2 ± 12.7

The Alizarin Red (2%) stained cells were incubated with 10% acetic acid for 30 minutes to release bound Alizarin Red into solution. The solution was neutralized with 10% ammonium hydroxide and the absorbance of Alizarin Red was measured at 450 nm using a microplate reader (n = 6). Data is expressed as absolute amounts according to a standard curve.

Conclusions

These data demonstrate the de-differentiation (reduction of mineralization genes) effect of FGF-2 likely plays a key role in osteoblast proliferation, the first stage of bone formation. Some have expressed concern that *ex-vivo* proliferation of human stem cells by a growth factor like FGF-2 might change the osteogenic characteristics of a pre-osteoblast; however others have shown that expansion of the population does not affect later osteogenic potential [35] of stem cells. Therefore, an expansion of osteoblast cells by FGF-2 might be an excellent strategy for first stage re-population of a critical defect since FGF-2

has the needed efficacy for promoting proliferation. These data also suggest that the final stage of bone repair is best accomplished with BMP-2 due to its promotion of differentiation and mineralization.

Acknowledgements

This work is supported by MHF's US Army Medical Research and Materiel Command US Army grant W81WH-07-1-0427, NASA grant NAG-2-1086 and in part by NASA grants NAG-2-1286, NCC2-1361 and the Department of Veterans Affairs Medical Research Service in support of MHF and this project. We thank Sandra Spurlock for the data plate reading and data analysis of Table 3. We would like to thank Tammy Chang for her thoughtful comments and suggestions during this work and Joe Meissler, Tara Candelario, Esmeralda Aguayo and Jesus Aguado for their thoughtful comments on the manuscript.

Author details

¹Department of Research, Veterans Affairs Medical Center, 4150 Clement Street, San Francisco, CA 94121, USA. ²Department of Medicine, University of California, 4150 Clement Street, San Francisco, CA 94121, USA. ³Department of Urology, University of California, 4150 Clement Street, San Francisco, CA 94121, USA. ⁴Hughes-Fulford Laboratory, Northern California Institute for Research and Education, 4150 Clement Street, San Francisco, CA 94121, USA.

Authors' contributions

MHF conceived the study, designed the study, directed the research and wrote the manuscript. C-FL made substantive intellectual contribution in the acquisition of data, analysis and has contributed to the manuscript. Both authors have read and approved the final manuscript.

Competing interests

The Department of Veterans Affairs has filed and owns a patent using some of the data found in this manuscript.

Received: 29 July 2010 Accepted: 9 February 2011

Published: 9 February 2011

References

1. Stein GS, Lian JB, Stein JL, Van Wijnen AJ, Montecino M: **Transcriptional control of osteoblast growth and differentiation.** *Physiol Rev* 1996, **76**:593-629.
2. Mundy GR, Chen D, Zhao M, Dallas S, Xu C, Harris S: **Growth regulatory factors and bone.** *Rev Endocr Metab Disord* 2001, **2**:105-115.
3. Canalis E, McCarthy TL, Centrella M: **Growth factors and cytokines in bone cell metabolism.** *Annu Rev Med* 1991, **42**:17-24.
4. Montero A, Okada Y, Tomita M, Ito M, Tsurukami H, Nakamura T, Doetschman T, Coffin JD, Hurley MM: **Disruption of the fibroblast growth factor-2 gene results in decreased bone mass and bone formation.** *J Clin Invest* 2000, **105**:1085-1093.
5. Burgess WH, Maciag T: **The heparin-binding (fibroblast) growth factor family of proteins.** *Annu Rev Biochem* 1989, **58**:575-606.
6. Santiago FS, Lowe HC, Day FL, Chésterman CN, Khachigian LM: **Early growth response factor-1 induction by injury is triggered by release and paracrine activation by fibroblast growth factor-2.** *Am J Pathol* 1999, **154**:937-944.
7. Grazul-Bilska AT, Johnson ML, Bilski JJ, Redmer DA, Reynolds LP, Abdullah A, Abdullah KM: **Wound healing: the role of growth factors.** *Drugs Today (Barc)* 2003, **787**:800.
8. Kodama HA, Amagai Y, Koyama H, Kasai S: **A new preadipose cell line derived from newborn mouse calvaria can promote the proliferation of pluripotent hemopoietic stem cells in vitro.** *J Cell Physiol* 1982, **112**:89-95.
9. Ibbotson KJ, Orcutt CM, Anglin AM, D'Souza SM: **Effects of transforming growth factors beta 1 and beta 2 on a mouse clonal, osteoblastlike cell line MC3T3-E1.** *J Bone Miner Res* 1989, **4**:37-45.
10. Katsuyama H, Otsuki T, Tomita M, Fukunaga M, Fukunaga T, Suzuki N, Saijoh K, Fushimi S, Sunami S: **Menaquinone-7 regulates the expressions of osteocalcin, OPG, RANKL and RANK in osteoblastic MC3T3E1 cells.** *Int J Mol Med* 2005, **15**:231-236.
11. Sierra OL, Cheng SL, Loewy AP, Charlton-Kachigian N, Towler DA: **MINT, the Msx2 interacting nuclear matrix target, enhances Runx2-dependent activation of the osteocalcin fibroblast growth factor response element.** *J Biol Chem* 2004, **279**:32913-32923.
12. Sudo H, Kodama HA, Amagai Y, Yamamoto S, Kasai S: **In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria.** *J Cell Biol* 1983, **96**:191-198.
13. Hatton JP, Pooran M, Li CF, Luzzio C, Hughes-Fulford M: **A short pulse of mechanical force induces gene expression and growth in MC3T3-E1 osteoblasts via an ERK 1/2 pathway.** *J Bone Miner Res* 2003, **18**:58-66.
14. Hughes-Fulford M, Rodenacker K, Jutting U: **Reduction of anabolic signals and alteration of osteoblast nuclear morphology in microgravity.** *J Cell Biochem* 2006, **99**:435-449.
15. Li CF, Hughes-Fulford M: **Fibroblast growth factor-2 is an immediate-early gene induced by mechanical stress in osteogenic cells.** *J Bone Miner Res* 2006, **21**:946-955.
16. Wiley MH, Feingold KR, Grunfeld C, Quesney-Huneeus V, Wu JM: **Evidence for cAMP-independent inhibition of S-phase DNA synthesis by prostaglandins.** *J Biol Chem* 1983, **258**:491-496.
17. Johnson RF, Mitchell CM, Giles WB, Bisits A, Zakar T: **Mechanisms regulating prostaglandin H2 synthase-2 mRNA level in the amnion and chorion during pregnancy.** *J Endocrinol* 2006, **188**:603-610.
18. Hughes-Fulford M, Gilbertson V: **Osteoblast fibronectin mRNA, protein synthesis, and matrix are unchanged after exposure to microgravity.** *Faseb J* 1999, **13**:S121-127.
19. Hughes-Fulford M, Tjandrawinata R, Fitzgerald J, Gasuad K, Gilbertson V: **Effects of microgravity on osteoblast growth.** *Gravit Space Biol Bull* 1998, **11**:51-60.
20. Tjandrawinata RR, Hughes-Fulford M: **Up-regulation of cyclooxygenase-2 by product-prostaglandin E2.** *Adv Exp Med Biol* 1997, **407**:163-170.
21. Hughes-Fulford M: **Prostaglandin regulation of gene expression and growth in normal and malignant tissues.** *Adv Exp Med Biol* 1997, **269**:278.
22. Hughes-Fulford M, Lewis ML: **Effects of microgravity on osteoblast growth activation.** *Exp Cell Res* 1996, **224**:103-109.
23. Stein GS, Lian JB, Stein JL, Van Wijnen AJ, Montecino M: **Transcriptional control of osteoblast growth and differentiation.** *Physiol Rev* 1996, **76**:593-629.
24. Hauschka PV, Chen TL, Mavrikos AE: **Polypeptide growth factors in bone matrix.** *Ciba Found Symp* 1988, **136**:207-225.
25. Tsuboi T, Mizutani S, Nakano M, Hirukawa K, Togari A: **Fgf-2 regulates enamel and dentine formation in mouse tooth germ.** *Calcif Tissue Int* 2003, **73**:496-501.
26. Gerstenfeld LC, Chipman SD, Kelly CM, Hodgins KJ, Lee DD, Landis WJ: **Collagen expression, ultrastructural assembly, and mineralization in cultures of chicken embryo osteoblasts.** *J Cell Biol* 1988, **106**:979-989.
27. Vary CP, Li V, Raouf A, Kitching R, Kola I, Franceschi C, Venanzoni M, Seth A: **Involvement of Ets transcription factors and targets in osteoblast differentiation and matrix mineralization.** *Exp Cell Res* 2000, **257**:213-222.
28. Chen D, Zhao M, Mundy GR: **Bone morphogenetic proteins.** *Growth Factors* 2004, **22**:233-241.
29. Mundy GR: **Regulation of bone formation by bone morphogenetic proteins and other growth factors.** *Clin Orthop Relat Res* 1996, **24**:28.
30. Laurent M, Martinerie C, Thibout H, Hoffman MP, Verrecchia F, Le Bouc Y, Mauviel A, Kleinman HK: **NOVH increases MMP3 expression and cell migration in glioblastoma cells via a PDGFR-alpha-dependent mechanism.** *Faseb J* 2003, **17**:1919-1921.
31. Koch H, Jadlovec JA, Campbell PG: **Insulin-like growth factor-I induces early osteoblast gene expression in human mesenchymal stem cells.** *Stem Cells Dev* 2005, **14**:621-631.
32. Wang Y, Nishida S, Sakata T, Elalieh HZ, Chang W, Halloran BP, Doty SB, Bikle DD: **Insulin-like growth factor-I is essential for embryonic bone development.** *Endocrinology* 2006, **147**:4753-4761.
33. Burghardt AJ, Wang Y, Elalieh H, Thibault X, Bikle D, Peyrin F, Majumdar S: **Evaluation of fetal bone structure and mineralization in IGF-I deficient mice using synchrotron radiation microtomography and Fourier transform infrared spectroscopy.** *Bone* 2007, **40**:160-168.
34. Betz OB, Betz VM, Nazarian A, Egermann M, Gerstenfeld LC, Einhorn TA, Vrahas MS, Bouxsein ML, Evans CH: **Delayed administration of adenoviral BMP-2 vector improves the formation of bone in osseous defects.** *Gene Ther* 2007, **14**:1039-1044.

35. Kulterer B, Friedl G, Jandrositz A, Sanchez-Cabo F, Prokesch A, Paar C, Scheideler M, Windhager R, Preisegger KH, Trajanoski Z: **Gene expression profiling of human mesenchymal stem cells derived from bone marrow during expansion and osteoblast differentiation.** *BMC Genomics* 2007, **8**:70.

doi:10.1186/1749-799X-6-8

Cite this article as: Hughes-Fulford and Li: The role of FGF-2 and BMP-2 in regulation of gene induction, cell proliferation and mineralization. *Journal of Orthopaedic Surgery and Research* 2011 **6**:8.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

