

CORRECTION

Open Access



Correction to: Achilles tenodesis for calcaneal insufficiency avulsion fractures associated with diabetes mellitus

Youngrak Choi¹, Young-woo Kwon¹, Young-suk Sim¹, Taeho Kim¹, Dayoung Song² and Soohyun Lee^{1*}

Correction

The original publication of this article [1] contained the wrong versions of Tables 1, 2 and 3. In this correction the updated tables are published. The original publication has been updated.

Author details

¹Department of Orthopedic Surgery, CHA Bundang Medical Center, CHA University, 16, Yatap-ro 65-beon-gil, Bundang-gu, Sungnam-si, Gyunggi-do 13497, Republic of Korea. ²School of Medicine, CHA University, 120, Haeryong-ro, Pocheon-si, Gyeonggi-do, Republic of Korea.

Received: 17 January 2018 Accepted: 15 February 2018

Published online: 02 March 2018

Reference

1. Choi Y, Kwon Y, Sim Y, et al. Achilles tenodesis for calcaneal insufficiency avulsion fractures associated with diabetes mellitus. *J Orthop Surg Res.* 2017;12:192. <https://doi.org/10.1186/s13018-017-0695-8>.

* Correspondence: chingize@gmail.com

¹Department of Orthopedic Surgery, CHA Bundang Medical Center, CHA University, 16, Yatap-ro 65-beon-gil, Bundang-gu, Sungnam-si, Gyunggi-do 13497, Republic of Korea

Full list of author information is available at the end of the article



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Table 1 Demographic Data of Patients

	No	Sex	Age	Period of follow-up (months)	PMHx	Duration of DM (years)	HgbA1c	Injury Mechanism	Pre-operative VAS
CIA fracture	1	F	63	14	DM	20	7.6	Ankle sprain	2
	2	M	55	15	DM ESRD	10	8.5	Walking	3
	3	F	48	20	DM	6	15.1	Walking	6
	4	M	51	18	DM	16	7.4	Climbing a hill	4
	5	F	48	18	DM ESRD	25	8.9	Climbing a hill	5
	6	M	43	16	DM	15	8.1	Walking	3
	7	M	79	20	DM ESRD	25	9.1	Walking	3

Table 2 The treatment process of Patients

	No	1 st operation	2 nd operation	3 rd operation
CIA fracture	1	ORIF ^a with screw (pulled out)	ORIF with TBW ^b (pulled out)	Achilles tenodesis and bone fragment resection
	2	Achilles tenodesis and bone fragment resection	Suture anchor removal because of pulled out failure	
	3	ORIF with screw (pulled out)	Achilles tenodesis and bone fragment resection	
	4	ORIF with screw (pulled out)	Achilles tenodesis and bone fragment resection	
	5	Achilles tenodesis and bone fragment resection		
	6	Achilles tenodesis and bone fragment resection		
	7	Achilles tenodesis and bone fragment resection		

^aOpen reduction and internal fixation^bTension band wiring**Table 3** Foot and Ankle Outcome Score (FAOS), Visual Analogue Scale (VAS) and single-heel rise test of Patients at final follow up

Group	No	FAOS					VAS	Single-heel rise test
		PAIN	SYMPTOMS	ADL ^a	SPORT&REC ^b	QOL ^c		
CIA fractures	1	77.8	82.1	85.3	75	81.3	3	+
	2	72.2	75.0	63.2	45	68.8	4	-
	3	88.9	89.3	85.3	75	87.5	3	+
	4	72.2	82.1	80.9	85	75	2	+
	5	77.8	82.1	77.9	80	81.3	3	+
	6	85.6	89.4	86.9	85	75	2	+
	7	89.5	86.9	83.8	85	76.4	1	+
	Mean	80.6	83.8	80.5	75.6	77.9	2.6	

^aActivities of daily living^bSport and recreation function^cFoot and ankle-related quality of life