CORRECTION Open Access



Correction to: Early versus late intramedullary nailing for traumatic femur fracture management: meta-analysis

Ayman El-Menyar^{1,2,3*}, Mohammed Muneer⁴, David Samson¹, Hassan Al-Thani⁵, Ahmad Alobaidi⁶, Paul Mussleman⁷ and Rifat Latifi⁸

Correction

Following the publication of this article [1], the authors reported that they had submitted an incorrect version of Figs. 2, 3 and 4. They apologize for this error and the correct versions of Figs. 2, 3 and 4 with captions have been included in this Correction. There is a typographical error in the following sentence: Two retrospective cohort studies reported results on any and all complications [23, 30]. The correct version of this sentence is: Two retrospective cohort studies reported results on any complication [24, 34].

Author details

¹Department of Surgery Clinical Research Unit, Westchester Medical Center Health Network, Valhalla, NY, USA. ²Trauma Surgery, Clinical Research, Hamad General Hospital, Doha, Qatar. ³Clinical Medicine, Weill Cornell Medical School, Doha, Qatar. ⁴Department of Surgery, Hamad General Hospital, Doha, Qatar. ⁵Department of Surgery, Trauma and Vascular Surgery, Hamad General Hospital, Doha, Qatar. ⁶Department of Surgery, Orthopedic Surgery, Al Wakrah Hospital, Doha, Qatar. ⁷Distributed eLibrary, Weill Cornell Medical School, Doha, Qatar. ⁸Department of Surgery, Westchester Medical Center Health Network and New York Medical College, Valhalla, NY, USA.

Received: 5 July 2018 Accepted: 6 July 2018 Published online: 24 July 2018

Reference

 El-Menyar, et al. Early versus late intramedullary nailing for traumatic femur fracture management: meta-analysis. J Orthop Surg Res. 2018;13:160. https://doi.org/10.1186/s13018-018-0856-4.


²Trauma Surgery, Clinical Research, Hamad General Hospital, Doha, Qatar Full list of author information is available at the end of the article



^{*} Correspondence: aymanco65@yahoo.com

¹Department of Surgery Clinical Research Unit, Westchester Medical Center Health Network, Valhalla, NY, USA

a) Pulmonary complications



b) Acute respiratory distress syndrome

	Early I	MN	Late II	ΝN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% C	I IV, Random, 95% CI
1, Alobaidi 2016	1	156	2	151	2.5%	0.48 [0.04, 5.36]	· · ·
22, Pape 2007	8	94	7	71	13.0%	0.85 [0.29, 2.47]	
24, Nahm 2011	7	408	4	84	9.4%	0.35 [0.10, 1.22]	
25, Bone 2004	1	88	6	90	3.2%	0.16 [0.02, 1.37]	· · ·
26, Charash 1994	2	105	3	33	4.4%	0.19 [0.03, 1.22]	
27, Starr 1998	0	14	1	18	1.4%	0.40 [0.02, 10.64]	
28, Boulanger 1997	4	125	4	27	7.0%	0.19 [0.04, 0.82]	
29, Brundage 2002	35	867	20	192	44.8%	0.36 [0.20, 0.64]	-
30, Harwood 2005	5	77	13	97	12.7%	0.45 [0.15, 1.32]	
31, O'Toole 2009	3	199	0	28	1.6%	1.02 [0.05, 20.17]	
Total (95% CI)		2133		791	100.0%	0.39 [0.26, 0.57]	•
Total events	66		60				
Heterogeneity: Tau ² =	0.00; Chi ²	= 4.79	df = 9 (F	= 0.85	5); I ² = 0%		0.01 0.1 1 10 100
Test for overall effect:	Z = 4.87 (P < 0.0	0001)				0.01 0.1 1 10 100 Favors early IMN Favors late IMN

c) Fat embolism

	Early I	MN	Late II	ΜN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
25, Bone 2004	0	88	2	90	9.2%	0.20 [0.01, 4.23]	-
26, Charash 1994	2	105	1	33	14.4%	0.62 [0.05, 7.08]	
29, Brundage 2002	10	867	3	192	50.5%	0.74 [0.20, 2.70]	
32, Al-Saflan 2012	2	90	3	105	26.0%	0.77 [0.13, 4.73]	-
Total (95% CI)		1150		420	100.0%	0.65 [0.26, 1.62]	-
Total events	14		9				
Heterogeneity: Tau ² =	0.00; Chi ²	= 0.64	, df = 3 (F		0.01 0.1 1 10 100		
Test for overall effect: 2	Z = 0.93 (P = 0.3	5)				Favors early IMN Favors late IMN

d) Pulmonary embolism


	Early If	MN	Late II	ΛN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% C	IV, Random, 95% CI
24, Nahm 2011	7	408	0	84	17.9%	3.16 [0.18, 55.81]	
25, Bone 2004	1	88	3	90	28.4%	0.33 [0.03, 3.27]	
26, Charash 1994	2	105	1	33	25.0%	0.62 [0.05, 7.08]	
27, Starr 1998	0	14	0	18		Not estimable	
29, Brundage 2002	3	867	1	192	28.7%	0.66 [0.07, 6.41]	-
Total (95% CI)		1468		399	100.0%	0.71 [0.21, 2.39]	
Total events	13		5				
Heterogeneity: Tau ² =	0.00; Chi ²	0.01 0.1 1 10 100					
Test for overall effect:	Z = 0.55 (F	P = 0.5	8)				Favors early IMN Favors late IMN

e) Pneumonia

	Early I	MN	Late II	MN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% C	IV, Random, 95% CI
22, Pape 2007	15	94	20	71	16.2%	0.48 [0.23, 1.03]	
24, Nahm 2011	40	408	19	84	21.4%	0.37 [0.20, 0.68]	
25, Bone 2004	1	88	3	90	2.5%	0.33 [0.03, 3.27]	
26, Charash 1994	13	105	15	33	12.7%	0.17 [0.07, 0.42]	
27, Starr 1998	1	14	10	18	2.6%	0.06 [0.01, 0.58]	
29, Brundage 2002	76	867	28	192	27.9%	0.56 [0.35, 0.90]	-
30, Harwood 2005	12	77	33	97	16.6%	0.36 [0.17, 0.75]	
Total (95% CI)		1653		585	100.0%	0.37 [0.26, 0.54]	•
Total events	158		128				
Heterogeneity: Tau ² = 0	0.07; Chi ²	= 8.68	df = 6 (P	= 0.19	6	0.01 0.1 1 10 100	
Test for overall effect: 2	Z = 5.20 (P < 0.0	0001)			Favors early IMN Favors late IMN	

Fig. 2 Forest plot of pulmonary complications. a Pulmonary complications. b Acute respiratory distress syndrome. c Fat embolism. d pulmonary embolism. e Pneumonia






b) Wound infection

	Early I	MN	Late II	ΝN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% C	I IV, Random, 95% CI
1, Alobaidi 2016	3	156	8	151	25.3%	0.35 [0.09, 1.35]	
21, Harvin 2012	9	1032	13	344	28.9%	0.22 [0.09, 0.53]	
24, Nahm 2011	3	408	1	84	18.3%	0.61 [0.06, 5.98]	
32, Al-Saflan 2012	14	90	5	105	27.5%	3.68 [1.27, 10.67]	_ -
Total (95% CI)		1686		684	100.0%	0.65 [0.15, 2.89]	
Total events	29		27				
Heterogeneity: Tau ² =	1.81; Chi ²	= 16.8	82%	0.01 0.1 1 10 100			
Test for overall effect:	Z = 0.56 (P = 0.5	7)				0.01 0.1 1 10 100 Favors early IMN Favors late IMN

c) Sepsis



d) Deep vein thrombosis

	Early I	MN	Late II	NN		Odds Ratio	Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
21, Harvin 2012	8	1032	10	344	38.1%	0.26 [0.10, 0.67]	
24, Nahm 2011	31	408	12	84	61.9%	0.49 [0.24, 1.01]	-
Total (95% CI)		1440		428	100.0%	0.39 [0.21, 0.71]	•
Total events	39		22				
Heterogeneity: Tau ² = Test for overall effect: 2				= 0.29	9); I² = 11%		0.01 0.1 1 10 100
							Favors early IMN Favors late IMN

Fig. 3 Forest plots of cutaneous, infectious and venous complications: a decubitus ulcers. b wound infection. c sepsis. d deep vein thrombosis




Fig. 4 Forest plots of other complications: a. Any complication. b multiorgan failure c. mortality