#### CORRECTION **Open Access**



# Correction: Aberrant expression of miR-33a-3p/IGF2 in postmenopausal osteoporosis patients and its role and mechanism in osteoporosis

Changxin Wang<sup>1</sup>, Jianfei Shen<sup>2</sup>, Wei Zhang<sup>3</sup>, Xiaoyu Wang<sup>2</sup>, Xiaohong Xu<sup>4</sup>, Xianghui Lu<sup>5</sup>, Dongbin Xu<sup>6</sup> and Lan Yao<sup>2\*</sup>

Correction: Journal of Orthopaedic Surgery and Research (2023) 18: 487

https://doi.org/10.1186/s13018-023-03883-6

In the sentence beginning 'Fifteen osteoporosis patients...' in this article, the incorrect text dual-energy X-ray bone densitometer (DEXA) should have read 'BMD was detected by a dual energy X-ray bone densitometer (DPX-NT, GE, WI, USA).

The updated sentence is given below and the changes have been highlighted in **bold typeface**.

The online version of the original article can be found at https://doi. ora/10.1186/s13018-023-03883-6.

\*Correspondence:

vaolan1982@163.com

Department of Orthopaedics, The Third Affiliated Hospital of Qiqihar Medical University, Qiqihar 161000, China

<sup>2</sup>Nuclear Medicine Department, The Third Affiliated Hospital of Qiqihar Medical University, No. 27 Taishun Street, Tiefeng Distric, Qigihar 161000 China

<sup>3</sup>Endocrine Department, The Third Affiliated Hospital of Qiqihar Medical University, Qigihar 161000, China

<sup>4</sup>Department of Clinical Laboratory, The Third Affiliated Hospital of Qiqihar Medical University, Qigihar 161000, China

 $^5$ Department of Gynaecology, The Third Affiliated Hospital of Qiqihar Medical University, Qiqihar 161000, China

<sup>6</sup>Qiqihar Medical University, Qiqihar 161000, China

### Materials and methods

#### **Patient characteristics**

Fifteen osteoporosis patients and healthy controls from the Third Affiliated Hospital of Qiqihar Medical University were included in the study. Written informed consent was obtained from all participants. Anthropometric characteristics, including age, BMI, and T-score of BMD, were recorded in the control and OP groups. The clinical data of osteoporosis patients and healthy volunteers were shown in Table 1. BMD was detected by a dual energy X-ray bone densitometer (DPX-NT, GE, WI, USA) according to WHO standards. This study was approved by the Ethics Committee of the Third Affiliated Hospital of Oigihar Medical University.

Published online: 12 September 2024

## Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© The Author(s) 2024. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will  $need to obtain permission directly from the copyright holder. To view a copy of this licence, visit \\ http://creativecommons.org/licenses/by/4.0/. The$ Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.