

RESEARCH ARTICLE

Open Access

Rabbiteye blueberry prevents osteoporosis in ovariectomized rats

Tao Li^{1†}, Shou-Mian Wu^{2†}, Zhi-Yuan Xu^{2*} and Sheng Ou-Yang^{2*}

Abstract

Objective: It has been forecasted that the rabbiteye blueberry could inhibit osteoporosis. However, the inhibition and prevention of osteoporosis via rabbiteye blueberry are still elusive. This study was aim to evaluate the anti-osteoporosis effects of rabbiteye blueberry in ovariectomized rats.

Methods: Thirty rats were randomly divided into three groups of ten rats each as follows: sham-operated group (SG), ovariectomized model control group (OMG), and ovariectomized rabbiteye blueberry treatment group (OBG). The blood mineral levels, the alkaline phosphatase (ALP) activity, and osteoprotegerin (OPG) level were determined. The expression analyses of type I collagen, integrin- β 1, and focal adhesion kinase (FAK) were performed. Besides, the bone mineral density (BMD) and bone histomorphometry (BH) were measured.

Results: The ALP activity in SG and OBG was significantly lower than that in OMG. For the OPG level, the significant increase of OPG level in OBG was indicated compared with the other groups. The mRNA expression levels of type I collagen, integrin- β 1, and FAK in OMG were significantly lower than those in other groups. The BMD in OMG were all significantly lower than those in SG and OBG. For BH, blueberry significantly improved the trabecular bone volume fraction, trabecular thickness, mean trabecular bone number, and bone formation rate, and decreased the trabecular separation, the percent of bone resorption perimeter, and mean osteoclast number in OBG compared with OMG.

Conclusions: The rabbiteye blueberries had an effective inhibition in bone resorption, bone loss, and reduction of bone strength of ovariectomized rats and could improve the BMD, osteogenic activity, and trabecular bone structure.

Keywords: Rabbiteye blueberry, Osteoporosis, Alkaline phosphatase activity, Focal adhesion kinase, Bone histomorphometry

Introduction

Osteoporosis is a systemic skeletal disease characterized by low bone mass and microarchitectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture [1]. In the USA, there are 10 million Americans with osteoporosis (8 million women and 2 million men) and another 34 million individuals with osteopenia, which resulted in a significant proportion of the population to be deemed 'at risk' [2].

The major current anti-osteoporotic therapy is pharmacologic treatment including bisphosphonates, estrogens,

selective estrogen receptor modulators, and parathyroid hormone [3]. In the past 5 years, additional drugs were likely to have been approved, such as the anti-RANK ligand monoclonal antibody [4] and cathepsin K inhibitors [5]. Most of the pharmacologic therapy strategies have been adopted that aim to inhibit excessive bone resorption and increase bone formation. There were also managements focused on non-pharmacologic treatments, such as a balanced diet and adequate calcium and vitamin D intake [6]. Folic acid [7], vitamin B₁₂ [8], carotenoid [9], and lycopene [10] have been reported effective in improving bone loss and relieving the symptoms of osteoporosis in postmenopausal women [11].

Rabbiteye blueberry (*Vaccinium ashei* Reade) is a highly heterozygous, cross-pollinating fruit species, and belongs

* Correspondence: guxin36y@hotmail.com; zhqinggan@163.com

[†]Equal contributors

²Department of Emergency, The Third Affiliated Hospital of Southern Medical University, Guangzhou 510630, China

Full list of author information is available at the end of the article

to the genus *Vaccinium* (*Vaccinium* spp.) and Ericaceae family. Recent reports showed that feeding a high-quality diet supplemented with blueberries to pre-pubertal rats or the rats at postnatal day 20 to 34 prevented ovariectomy-induced bone loss in adult life [12]. The molecular mechanisms underlying these effects may involve the increase of myosin production and reduction of mesenchymal stromal cell senescence. However, the effect of blueberry diet on inhibiting ovariectomy-induced osteoporosis and its mechanisms still need research.

In order to provide more evidence and gain a better understanding on the association between the active ingredients of rabbiteye blueberries and the inhibition of ovariectomy-induced osteoporosis, *in vivo* experiments on ovariectomized rat models have been conducted in this study. The blood mineral levels, the activity of alkaline phosphatase (ALP), and osteoprotegerin (OPG) level were determined to evaluate the effect of rabbiteye blueberry to osteoporosis. The expression analysis of type I collagen, integrin- β 1, and focal adhesion kinase (FAK) was performed to explore the influence of rabbiteye blueberries on the osteoporosis-related integrin pathway. Further measurements of bone mineral density (BMD) and bone histomorphometry (BH) were performed to observe bone microstructure and dynamic changes of bone metabolism in osteoporosis.

Materials and methods

Experimental animals

Our experimental research on animals followed internationally recognized guidelines. The Animal Ethics Committee of Shanghai Jiao Tong University School of Medicine approved the study (reference number 2010–0018).

A total of 30 normal 8-week-old female Sprague–Dawley rats (160 ± 15 g) were provided by the Laboratory of Animal Science Department of Shanghai Jiao Tong University School of Medicine. The 30 rats were housed in ten cages with temperature of $21^\circ\text{C} \pm 1^\circ\text{C}$, relative humidity of 60%, ambient noise <50 dB, and regular 12-h light cycle. All rats were feed with conventional food and free access to water.

Operative techniques

The 30 rats were randomly divided into three groups of ten rats each as follows: sham-operated group (SG), ovariectomized model control group (OMG), and ovariectomized blueberry treatment group (OBG). All rats were anesthetized before operation by intraperitoneal injection with 1% (volume fraction) pentobarbital sodium at a dose of 40 mg/kg. The operation begun through a 1-cm longitudinal incision at the back, 1 cm from the spine and 2 cm above the posterior iliac crest. Bilateral oophorectomy was performed on the rats of OMG and OBG, while same weights of greater omentums were

removed instead of ovary for the rats of SG. Finally, the wound was closed in layers after stump ligation.

Rabbiteye blueberry administration

After the operation, penicillin was intramuscularly injected to all the rats for three consecutive days. One week later, the rats in the OBG were administered with freeze-dried blueberry powder water solution (10%, *w/w*) for 12 consecutive weeks by gavage, while the rats in the SG and OMG were administered with equal volume of double distilled water by gavage. The freeze-dried blueberry powder (stored at -20°C) was provided by the Institute of Botany, Chinese Academy of Sciences, Beijing, China.

Determination of the levels of serum minerals, ALP, and OPG

After the 12-week administration, the rats were anesthetized by intraperitoneal injection of 20% (*w/v*) urethane solution at a dose of 1 mg/kg. The blood samples were collected from the abdominal aorta and then the serum was obtained by centrifugation at a low temperature of 4°C . The levels of the serum minerals were determined by using an automatic biochemical analyzer. The ALP activity was determined using an Alkaline Phosphatase Opt Kit (Roche Molecular Biochemical, Indianapolis, IN, USA). The OPG level in serum was measured by ELISA using the Biomedical rat OPG ELISA kit (Biomedical, Wien, Austria). All measurements were performed according to the manufacturer's instructions.

Total RNA extraction and reverse transcription of RNA

After being fasted for 24 h, all the rats were sacrificed. Then, the third lumbar vertebra without surrounding soft tissues was quickly collected at once. After homogenized treatment, the total RNA was extracted by using Trizol Reagent (Shanghai Jianglai Biotechnology Co., Ltd., Shanghai, China) according to the manufacturer's recommendations. The quantity and quality of RNA were determined by UV absorbance at 260-, 280-, and 320-nm wavelengths using NanoDrop ND-1000 (Thermo Fisher Scientific Inc., Waltham, MA, USA). Complementary DNA (cDNA) was synthesized from RNA using Reverse-transcription kit (Life Technologies Company, Beijing, China) with Superscript II Reverse Transcriptase (Life Technologies Inc., CA, USA) according to the manufacturer's recommendations.

Detection of expression of type I collagen, integrin- β 1, and focal adhesion kinase

The real-time quantitative polymerase chain reaction (PCR) was carried out by CFX96 Real-Time PCR instrument (Bio-Rad, Hercules, CA, USA). SYBR Green I was provided by Shanghai Life Technology Company

Table 1 Primers of collagen-1, integrin-β1, FAK, and β-actin

Gene name	Forward primer	Reverse primer
<i>Collagen-1</i>	ATCAAGGTCACCTGCAACAT	CAGGATCAAACCTTCGCTT
<i>Integrin-β1</i>	CCCTTTCCTCAGAAGTCATTTTG	TTCGCTTTGGCATTACATTAC
<i>FAK</i>	CGGGATCCGCATCAGCATA	CCCAAGCTTTTAGTGCCCTG
<i>β-actin</i>	TCT TCC AGC CTT CCT TCC TG	TAG AGC CAC CAA TCC ACA CA

Collagen-1 encodes type I collagen, *integrin-β1* encodes integrin-β1, and *FAK* encodes focal adhesion kinase.

(Shanghai, China). Primers of *collagen-1* (encoding type I collagen), *integrin-β1* (encoding integrin-β1), *FAK* (encoding focal adhesion kinase), and *β-actin* were shown in Table 1. The real-time quantitative PCR was carried out under the following conditions: preheating at 95°C for 5 min, followed by 45 cycles of denaturation at 95°C for 1 min, annealing at 54°C for 1 min, and extension at 74°C for 2 min. The PCR amplification was ended by a final extension cycle at 72°C for 10 min. After obtaining the amplification curve and the CT values, the dissolve curve was analyzed using the following setting: 95°C for 15 s, 60°C for 1 min, and 95°C for 15 s. The relative expression of each biomarker was represented as the ratio of the copy number of each sample and *β-actin*.

Detection of BMD

The left femur BMD was measured by using EXA-3000 dual-energy X-ray BMD measure instrument (MEDILINK, Maugeio, France). Based on actual length measured by the measuring instrument, the left femur BMD was divided into three parts: the proximal BMD (pBMD) in the upper portion including one fourth of the femur, the middle (mBMD) in middle portion including one half of the femur, and the distance (dBMD) in the lower end portion including one fourth of the femur.

Detection of histomorphometry

The complete right femur and tibia were used to prepare bone specimens for analyzing BH. The bone specimens were stained by von Kossa and Giemsa before they were observed and imaged by fluorescent/optical microscope (Leica DM4000B, Stuttgart, Germany). Randomly selected images in the inner, middle, and outer of the specimen were saved.

Static and dynamic histomorphometric measurements of the secondary cancellous bone located within 4.0 mm distal to the growth plate-metaphyseal were performed. The specimens stained with von Kossa were used for histology observation, while the specimens stained with Giemsa were used to observe osteoid. The data of BH were analyzed by using a color image analysis software of Leica.

Statistical analysis

The SPSS 12.0 software (<http://www-01.ibm.com/software/analytics/spss/>) was used for the statistical analysis. The results were all expressed as mean ± SEM. The analysis of variance and multiple comparison tests were performed with *p* value less than 0.05, which was considered as significant.

Results

Serum mineral levels

The serum minerals were monitored for changes that may affect bone mineralization. The levels of serum minerals (Ca, P, and Mg) in each group were shown in percentages (Table 2). The results indicated that there was no significant difference in the levels of serum minerals (Ca, P, and Mg) among the three groups with *p* value of more than 0.05.

OPG level and ALP activity

The ALP activity in SG and OBG was significantly lower than that in OMG (SG vs. OMG: *p* = 0.035; OBG vs. OMG: *p* = 0.041), and there was no significant difference between SG and OBG. For the OPG level in the serum, the significant increase of OPG level in OBG was indicated when compared with the other groups (OBG vs. SG: *p* = 0.029). The OPG level in SG was significantly lower than that in OMG (Figure 1).

The mRNA expression levels of type I collagen, integrin-β1, and FAK

Compared with the OMG, the mRNA expression levels of type I collagen, integrin-β1, and FAK were significantly higher in the OBG and SG (type I collagen: SG vs. OMG, *p* = 0.005; OBG vs. OMG, *p* = 0.001; integrin-β1: SG vs.

Table 2 The levels of serum minerals (Ca, P, and Mg) in each group

Name	OMG	SG	OBG
Ca (mmol/L)	2.20 ± 0.16	2.34 ± 0.08	2.22 ± 0.21
P (mmol/L)	1.62 ± 0.24	1.71 ± 0.17	1.82 ± 0.14
Mg (mmol/L)	0.63 ± 0.11	0.52 ± 0.08	0.53 ± 0.12

OMG ovariectomized model control group, SG sham-operated group, OBG ovariectomized rabbit eye blueberry treatment group.




Figure 1 The alkaline phosphatase (ALP) activity and osteoprotegerin (OPG) level in each group ($n = 10$). The activity (right image) and level (left image) were compared with those in the ovariectomized model control group ($*p < 0.05$, $**p < 0.01$) and compared with those in the sham-operated group ($\#p < 0.05$). Model: ovariectomized model control group (OMG); sham: sham-operated group (SG); blueberry: ovariectomized blueberry treatment group (OBG).

OMG, $p = 0.008$; OBG vs. OMG, $p = 0.001$; FAK: SG vs. OMG, $p = 0.008$; OBG vs. OMG, $p = 0.001$). Moreover, their mRNA expression levels in OBG were significantly increased compared with those in SG (type I collagen: OBG vs. SG, $p = 0.001$; integrin- $\beta 1$: OBG vs. SG, $p = 0.001$; FAK: OBG vs. SG, $p = 0.001$) (Figure 2).

Comparison of BMD

The total BMD, pBMD, mBMD, and dBMD in OMG were all significantly lower than those in SG (total BMD: $p = 0.012$; pBMD: $p = 0.025$; mBMD: $p = 0.016$; dBMD: $p = 0.021$). Compared with the OMG, the total BMD, pBMD, mBMD, and dBMD were all significantly increased in OBG (total BMD: $p = 0.028$; pBMD: $p = 0.036$; mBMD: $p = 0.045$; dBMD: $p = 0.042$) (Figure 3).

Bone histomorphology

The statistical analysis for the results of histomorphometric measurements indicated that the static and dynamic parameters of BH in SG and OBG presented significant differences compared with the OMG. The trabecular bone volume fraction (trabecular bone volume (BV)/total tissue volume (TV)), trabecular thickness (Tb. Th), and mean trabecular bone number (Tb. N) in OBG and SG were significant higher than those in OMG ($p < 0.01$). The trabecular separation (Tb. Sp), the percent of bone resorption perimeter (ES/BS), and the mean osteoclast number (Oc. N) in OBG and SG were significantly lower than those in OMG ($p < 0.05$). The osteoid perimeter (O. Pm) in SG was significantly lower than that in OMG ($p < 0.05$), while there was no significant difference in the osteoid perimeter between OBG and OMG. The fluorescent tag circumference ratio (L. Pm) was significantly lower in SG than in OMG ($p < 0.01$) but significantly higher in OBG than in OMG ($p < 0.05$). The bone formation rate (BFR/BV) in OBG and SG was higher than that in OMG ($p < 0.01$) (Table 3).

Discussion

Osteoporosis is a systemic skeletal disease that is often accompanied with a decrease of bone mass and micro-architectural deterioration of bone tissue, which can lead to an enhanced bone fragility and consequent increase in fracture risk [13]. The risk is greater in postmenopausal women due to deficiency of estrogen [14], which was considered the seminal mechanism of osteoporosis in both women and men [15]. Thus, a further understanding on osteoporosis pathogenesis, combined with the development of treatments with proven efficacy, will increase demand for the more effective management of patients with osteoporosis [16]. The lyophilized rabbiteye blueberries have been reported to have antioxidant activity [17] and can affect lipid metabolism [18]. In addition, it was reported that GTP supplementation plus





Figure 2 mRNA expression levels of type I collagen, integrin- $\beta 1$, and FAK in each group ($n = 10$). The levels were compared with those in the ovariectomized model control group ($*p < 0.01$) and compared with those in the sham-operated group ($\#p < 0.01$). Model: ovariectomized model control group (OMG); sham: sham-operated group (SG); blueberry: ovariectomized blueberry treatment group (OBG).



alphacalcidol administration increased bone mass via a decrease of oxidative stress [19]. Therefore, its antioxidant activity may be one of the mechanisms for preventing osteoporosis. However, there are few studies focusing on the preventive effects of rabbiteye blueberries on postmenopausal osteoporosis which is the most common cause of age-related bone loss.

Ovariectomy-induced bone loss and postmenopausal bone loss in the rat share many similar characteristics [20]. Thus, the ovariectomy-induced osteoporosis in rat was used as a model for postmenopausal osteoporosis in humans in this study. Bone mineral mainly includes Ca, P, and Mg, and bone metabolism can be indirectly studied through their levels in the serum. In this study, their levels presented no significant difference among the

three groups. The reason may be that the loss of bone mineral presents as a slow process due to a more active bone resorption than bone formation in the ovariectomized rats.

The ALP activity in the OMG presented a sharp rise compared with those in the other groups and no significant difference between OBG and SG, suggesting that the rabbiteye blueberry diet played a bone-protective role in the ovariectomized rat model. Moreover, the OPG level in OBG was significantly increased compared with those in the OMG and SG. It is well known that the balance between bone formation and bone resorption is essential for bone mass and bone microstructure [21]. ALP is assessed as a marker of bone formation [22], while OPG is identified as an inhibitor of osteoclastogenesis, osteoclast differentiation, and activation both *in vitro* and *in vivo* [23]. Significantly lower ALP activity and higher OPG level in OBG than in OMG were detected, revealing that rabbiteye blueberry can prevent osteoblast development by increasing the OPG level and can prevent osteoporosis by inhibiting the ALP activity.

Integrins are heterodimeric transmembrane proteins consisting of α and β subunits that functionally serve as both adhesive and signaling receptors [24]. Integrin- β 1 is a major receptor of matrix glycoproteins in type I collagen-induced osteoblastic differentiation of bone marrow cells [25]. Type I collagen interacts with integrin- β 1 receptor on the cell membrane, mediates extracellular signals into cells, and activates the FAK signal pathway. FAK is the immediate transducer in integrin- β 1-dependent adhesion of osteoblasts to bone matrices [26]. In this study, we observed that the mRNAs of type I collagen, integrin- β 1, and FAK in OMG expressed less than those in SG and OBG, and they expressed more in OBG than in SG. Ovariectomy could induce estrogen deficiency and increase

Table 3 Results of bone histomorphometry in each group

	OMG	SG	p value	OBG	p value
BV/TV (%)	16.38 ± 5.69	29.47 ± 6.52*	0.001	22.39 ± 5.26*	0.005
Tb. Th (μm)	48.96 ± 8.64	85.36 ± 9.35*	0.001	72.96 ± 7.37*	0.002
Tb. N (L/mm)	1.69 ± 0.16	2.05 ± 0.16*	0.005	1.89 ± 0.11*	0.008
Tb. Sp (μm)	763.85 ± 138.24	369.42 ± 102.34*	0.001	587.56 ± 185.21*	0.008
ES/BS (%)	18.27 ± 4.82	9.50 ± 4.42*	0.007	14.27 ± 4.23**	0.044
Oc. N (L/mm ²)	0.26 ± 0.08	0.10 ± 0.06*	0.001	0.20 ± 0.05**	0.039
O. Pm (%)	30.35 ± 6.32	18.84 ± 6.47**	0.030	31.13 ± 5.08	-
L. Pm (%)	15.03 ± 1.26	8.63 ± 1.09*	0.001	16.98 ± 3.25**	0.041
BFR/BV (%/year)	36.85 ± 6.23	58.27 ± 7.03*	0.002	61.23 ± 13.91*	0.001

*p < 0.01 and **p < 0.05 compared with the OMG. OMG ovariectomized model control group, SG sham-operated group, OBG ovariectomized rabbiteye blueberry treatment group, BV/TV trabecular bone volume fraction (trabecular bone volume (BV)/total tissue volume (TV)), Tb. Th trabecular thickness, Tb. N mean trabecular bone number, Tb. Sp trabecular separation, ES/BS the percent of bone resorption perimeter, Oc. N mean osteoclast number, O. Pm osteoid perimeter, L. Pm fluorescent tag circumference ratio, BFR/BV bone formation rate.

bone resorption which led to accelerated collagen degradation and reduced collagen synthesis, especially type I collagen [27,28]; then, the expression of integrin- β 1 and FAK may be affected. The reason may be that the mRNA expression levels of type I collagen, integrin- β 1, and FAK in the model rats were decreased in OMG. Meanwhile, the mRNA expression levels of type I collagen, integrin- β 1, and FAK in OBG were significantly higher than those in the other groups, indicating that rabbiteye blueberry played a positive role in preventing osteoporosis. The integrin- β 1/FAK signaling pathway may be repaired by active ingredients in blueberry to induce the increase of osteogenic activity.

Bone strength is determined by bone mass and bone quality. Bone mass is estimated in clinical practice by measuring the BMD. Menopause is the largest contributor to decreasing bone mineral density due to the significant decrease of estrogen [29]. The total BMD, pBMD, mBMD, and dBMD in OMG were all significantly lower than those in SG, and blueberry increased the BMD in OBG compared with OMG. This indicated that blueberry could effectively improve BMD of ovariectomized rats. Compared to BMD, BH can reflect the bone quality including bone microstructure and dynamic changes of bone metabolism. Trabecular bone belongs to cancellous bone, which is more sensitive to osteopenia than the cortical bone [30]. The trabecular architecture becomes more rod-like and anisotropic in osteoporotic and aging trabecular bone [31,32], even completely disappears in postmenopausal women and ovariectomized rats [33]. In this study, rabbiteye blueberries significantly improved the trabecular bone volume fraction, trabecular thickness, and mean trabecular bone number, and decrease the trabecular separation in OBG compared with OMG. Thus, the active ingredients in blueberry could prevent the loss of trabecular bone in osteoporosis.

It was reported that the decrease of estrogen enhanced the osteoclast development and increased the number of osteoclasts in the trabecular bone [34]. According to the results of this study, the percent of bone resorption perimeter and mean osteoclast number in OBG and SG were significantly lower than those in OMG, and the bone formation rate in OBG and SG were higher than that in OMG, indicating that the blueberry effectively inhibited the development of osteoclast and bone resorption and promoted bone formation. However, the osteoid perimeter and the fluorescent tag circumference ratio in SG were significantly lower than that in OMG, and blueberry did not inhibit their increase in OBG. The active ingredients in blueberries were complex. There may be some ingredients that can affect the osteoid perimeter and the fluorescent tag circumference ratio. More studies were required to explore the active ingredients in blueberries for preventing osteoporosis.

Conclusion

In conclusion, we have validated the inhibition of rabbiteye blueberry on ovariectomy-induced osteoporosis *in vivo*. The results showed that the rabbiteye blueberries had an effective inhibition in bone resorption, bone loss, and reduction of bone strength of ovariectomized rats, and could improve the BMD, osteogenic activity, and trabecular bone structure. Therefore, we conclude that blueberry might prevent and inhibit ovariectomy-induced osteoporosis, which provides new basis for the clinical treatment of osteoporosis. We showed, for the first time, that rabbiteye blueberries have great potential to accurately, reliably, and safely improve osteoporosis in a rat model of ovariectomy-induced osteoporosis.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

TL participated in the design of this study and performed the statistical analysis. SW carried out the study, together with SO, collected important background information, and drafted the manuscript. ZX conceived this study, participated in the design, and helped draft the manuscript. All authors read and approved the final manuscript.

Author details

¹Department of Orthopedics, The Third Affiliated Hospital of Southern Medical University (Academy of Orthopedics), Guangzhou, Guangdong Province 510630, China. ²Department of Emergency, The Third Affiliated Hospital of Southern Medical University, Guangzhou 510630, China.

Received: 20 January 2014 Accepted: 20 June 2014

Published: 8 August 2014

References

1. WHO Scientific Group on Prevention and Management of Osteoporosis: Prevention and management of osteoporosis: report of a WHO Scientific Group. *World Health Organ Tech Rep Ser* 2003, **921**:1–164.
2. Brömme D, Lecaille F: Cathepsin K inhibitors for osteoporosis and potential off-target effects. *Expert Opin Investig Drugs* 2009, **18**:585–600.
3. Tafaro L, Nati G, Leoni E, Baldini R, Cattaruzza M, Mei M, Falaschi P: Adherence to anti-osteoporotic therapies: role and determinants of "spot therapy". *Osteoporos Int* 2013, **24**(8):2319–2323.
4. Taylor K, Middlefell L, Mizen K: Osteonecrosis of the jaws induced by anti-RANK ligand therapy. *Br J Oral Maxillofac Surg* 2010, **48**(3):221–223.
5. Boonen S, Rosenberg E, Claessens F, Vanderschueren D, Papapoulos S: Inhibition of cathepsin K for treatment of osteoporosis. *Curr Osteoporos Rep* 2012, **10**(1):73–79.
6. McDermott MT: *Osteoporosis 2011: Compromised Bone Strength Predisposing to Increased Risk of Fragility and Fractures*. Aurora: University of Colorado Hospital; 2011.
7. Armitage JM, Bowman L, Clarke RJ, Wallendzus K, Bulbulia R, Rahimi K, Haynes R, Parish S, Sleight P, Peto R: Effects of homocysteine-lowering with folic acid plus vitamin B12 vs placebo on mortality and major morbidity in myocardial infarction survivors. *JAMA* 2010, **303**(24):2486–2494.
8. Baines M, Kredan M, Usher J, Davison A, Higgins G, Taylor W, West C, Fraser W, Ranganath L: The association of homocysteine and its determinants MTHFR genotype, folate, vitamin B12 and vitamin B6 with bone mineral density in postmenopausal British women. *Bone* 2007, **40**(3):730–736.
9. Sugiura M, Nakamura M, Ogawa K, Ikoma Y, Yano M: High serum carotenoids associated with lower risk for bone loss and osteoporosis in post-menopausal Japanese female subjects: prospective cohort study. *PLoS One* 2012, **7**(12):e52643.
10. Sahni S, Hannan MT, Blumberg J, Cupples LA, Kiel DP, Tucker KL: Protective effect of total carotenoid and lycopene intake on the risk of hip fracture: a 17-year follow-up from the Framingham Osteoporosis Study. *J Bone Miner Res* 2009, **24**(6):1086–1094.

11. Sugiura M, Nakamura M, Ogawa K, Ikoma Y, Ando F, Shimokata H, Yano M: **Dietary patterns of antioxidant vitamin and carotenoid intake associated with bone mineral density: findings from post-menopausal Japanese female subjects.** *Osteoporos Int* 2011, **22**(1):143–152.
12. Zhang J, Lazarenko OP, Blackburn ML, Shankar K, Badger TM, Ronis MJ, Chen J-R: **Feeding blueberry diets in early life prevent senescence of osteoblasts and bone loss in ovariectomized adult female rats.** *PLoS One* 2011, **6**(9):e24486.
13. Cooper C: **Osteoporosis: disease severity and consequent fracture management.** *Osteoporos Int* 2010, **21**(2):425–429.
14. Osako MK, Nakagami H, Koibuchi N, Shimizu H, Nakagami F, Koriyama H, Shimamura M, Miyake T, Rakugi H, Morishita R: **Estrogen inhibits vascular calcification via vascular RANKL system: common mechanism of osteoporosis and vascular calcification.** *Circ Res* 2010, **107**(4):466.
15. Manolagas SC: **From estrogen-centric to aging and oxidative stress: a revised perspective of the pathogenesis of osteoporosis.** *Endocr Rev* 2010, **31**(3):266–300.
16. Kanis J, Black D, Cooper C, Dargent P, Dawson-Hughes B, De Laet C, Delmas P, Eisman J, Johnell O, Jonsson B: **A new approach to the development of assessment guidelines for osteoporosis.** *Osteoporos Int* 2002, **13**(7):527–536.
17. Ramirez MR, Guterres L, Dickel OE, de Castro MR, Henriques AT, de Souza MM, Barros DM: **Preliminary studies on the antinociceptive activity of Vaccinium ashei berry in experimental animal models.** *J Med Food* 2010, **13**(2):336–342.
18. Inoue N, Nagao K, Nomura S, Shirouchi B, Inafuku M, Hirabaru H, Nakahara N, Nishizono S, Tanaka T, Yanagita T: **Effect of Vaccinium ashei Reade leaf extracts on lipid metabolism in obese OLETF rats.** *Biosci Biotechnol Biochem* 2011, **75**(12):2304–2308.
19. Shen C-L, Yeh J, Cao J, Tatum O, Dagda R, Wang J-S: **Synergistic effects of green tea polyphenols and alphacalcidol on chronic inflammation-induced bone loss in female rats.** *Osteoporos Int* 2010, **21**(11):1841–1852.
20. Shuid AN, Ping LL, Muhammad N, Mohamed N, Soelaiman IN: **The effects of Labisia pumila var. alata on bone markers and bone calcium in a rat model of post-menopausal osteoporosis.** *J Ethnopharmacol* 2011, **133**(2):538–542.
21. Gennari L, Merlotti D, Martini G, Gonnelli S, Franci B, Campagna S, Lucani B, Dal Canto N, Valenti R, Gennari C, Nuti R: **Longitudinal association between sex hormone levels, bone loss, and bone turnover in elderly men.** *J Clin Endocrinol Metab* 2003, **88**(11):5327–5333.
22. Otto F, Lubbert M, Stock M: **Upstream and downstream targets of RUNX proteins.** *J Cell Biochem* 2003, **89**(1):9–18.
23. Bucay N, Sarosi I, Dunstan CR, Morony S, Tarpley J, Capparelli C, Scully S, Tan HL, Xu W, Lacey DL, Boyle WJ, Simonet WS: **Osteoprotegerin-deficient mice develop early onset osteoporosis and arterial calcification.** *Genes Dev* 1998, **12**(9):1260–1268.
24. Shekaran A, Garcia AJ: **Extracellular matrix-mimetic adhesive biomaterials for bone repair.** *J Biomed Mater Res A* 2011, **96**(1):261–272.
25. Mizuno M, Fujisawa R, Kuboki Y: **Type I collagen-induced osteoblastic differentiation of bone-marrow cells mediated by collagen- $\alpha 2 \beta 1$ integrin interaction.** *J Cell Physiol* 2000, **184**(2):207–213.
26. Reyes CD, Garcia AJ: **$\alpha 2 \beta 1$ integrin-specific collagen-mimetic surfaces supporting osteoblastic differentiation.** *J Biomed Mater Res A* 2004, **69**(4):591–600.
27. Garnero P, Gineyts E, Riou JP, Delmas PD: **Assessment of bone resorption with a new marker of collagen degradation in patients with metabolic bone disease.** *J Clin Endocrinol Metab* 1994, **79**(3):780–785.
28. Risteli J, Niemi S, Elomaa I, Risteli L: **Bone resorption assay based on a peptide liberated during type I collagen degradation.** *J Bone Miner Res* 1991, **6**(Suppl 1):S251.
29. Riggs BL, Khosla S, Melton LJ III: **Sex steroids and the construction and conservation of the adult skeleton.** *Endocr Rev* 2002, **23**(3):279–302.
30. Francis DA, Millis DL, Head LL: **Bone and lean tissue changes following cranial cruciate ligament transection and stifle stabilization.** *J Am Anim Hosp Assoc* 2006, **42**(2):127–135.
31. David V, Laroche N, Boudignon B, Lafage-Proust MH, Alexandre C, Rueggsegger P, Vico L: **Noninvasive in vivo monitoring of bone architecture alterations in hindlimb-unloaded female rats using novel three-dimensional microcomputed tomography.** *J Bone Miner Res* 2003, **18**(9):1622–1631.
32. Ding M, Hvid I: **Quantification of age-related changes in the structure model type and trabecular thickness of human tibial cancellous bone.** *Bone* 2000, **26**(3):291–295.
33. Burguera B, Hofbauer LC, Thomas T, Gori F, Evans GL, Khosla S, Riggs BL, Turner RT: **Leptin reduces ovariectomy-induced bone loss in rats.** *Endocrinology* 2001, **142**(8):3546–3553.
34. Jilka RL, Hangoc G, Girasole G, Passeri G, Williams DC, Abrams JS, Boyce B, Broxmeyer H, Manolagas SC: **Increased osteoclast development after estrogen loss: mediation by interleukin-6.** *Science* 1992, **257**(5066):88–91.

doi:10.1186/s13018-014-0056-9

Cite this article as: Li et al.: Rabbiteye blueberry prevents osteoporosis in ovariectomized rats. *Journal of Orthopaedic Surgery and Research* 2014 9:56.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

